


Rabies Post-Exposure Prophylaxis for Hospitals

Please complete this form and fax to 519-539-5059

Reporting Hospital: _____

Ordering Physician: _____

Patient Name: _____



Patient Date of Birth: _____ Patient Weight (kg): _____

Rabies Immune Globulin (Rablg) *

Rablg **must** be calculated* for each client. The recommended dose is 20 IU/kg body weight for all age groups, given on the first day of initiation of therapy. Immune globulin is supplied in 150 IU/ml (2 ml) or 300 IU/ml (1 ml) vials. Check the vial label carefully when calculating the dosage (see reverse).

Examples for a client weighing 125 kg

- 300 IU/ml formulation of Rablg in 1 ml vials is calculated as follows:
(20 IU/kg x 125 kg ÷ 300 IU/ml = **8.3 ml**)
- 150 IU/ml formulation of Rablg in 2 ml vials is calculated as follows:
(20 IU/kg x 125 kg ÷ 150 IU/ml = **16.7 ml**)

Please select one: 300 IU/ml or 150 IU/ml

Day 0: _____ volume of Rablg in ml administered: _____
(date)

Rablg Lot # _____ Expiry Date: _____

Rabies Vaccine **

Day 0: _____ Lot # _____ Expiry Date: _____
(date)

Day 3: _____ Lot # _____ Expiry Date: _____
(date)

Day 7: _____ Lot # _____ Expiry Date: _____
(date)

Day 14: _____ Lot # _____ Expiry Date: _____
(date)

For previously **unimmunized immunocompromised** persons (including those taking corticosteroids, chloroquine or other antimalarials) receive a 5th dose of vaccine.

Day 28: _____ Lot # _____ Expiry Date: _____
(date)

NOTE

* If anatomically feasible, the full dose of **Rabies Immune Globulin** should be thoroughly infiltrated into the wound and surrounding area. Any remaining volume should be injected intramuscularly using a separate syringe at a site distant from administration of rabies vaccine, unless directed differently from your Public Health Department.

** To get the best antibody response for those ≥ one year, **Rabies Vaccine** should be given intramuscularly into the deltoid muscle. Infants less than one year, the vastus lateralis muscle (anterolateral thigh) is the preferred site. **DO NOT** give subcutaneously, and **NEVER** in the gluteal muscle.

2019 Changes to Formulations of Rablg Available in Ontario

As of July 2019, Rablg will be available in one of two formulations:

- 2 mL vials containing 150 IU/ml; or
- 1 mL vials containing 300 IU/ml.

Ensure that the appropriate formula specific to the Rablg formulation being provided to the physician is used to calculate the dose required for the individual to receive Rablg, and use **Table 1** to determine how many vials to dispense:

For 150 IU/ml Rablg in 2 ml vials:

- $20 \text{ IU/kg} \times (\text{client wt in kg}) \div 150 \text{ IU/mL} = \text{dose in mL}$
dose in mL \div 2 mL/vial = # of vials to order
- $9.09 \text{ IU/lb} \times (\text{client wt in lb}) \div 150 \text{ IU/mL} = \text{dose in mL}$
dose in mL \div 2 mL/vial = # of vials to order

For 300 IU/ml Rablg in 1 ml vials:

- $20 \text{ IU/kg} \times (\text{client wt in kg}) \div 300 \text{ IU/mL} = \text{dose in mL}$
dose in mL \div 1 mL/vial = # of vials to order
- $9.09 \text{ IU/lb} \times (\text{client wt in lb}) \div 300 \text{ IU/mL} = \text{dose in mL}$
dose in mL \div 1 mL/vial = # of vials to order

Note that while the dose in mL to be administered will be different depending on which formulation of Rablg is being used, the number of vials to be dispensed will still be the same.

Table 1: Number of Vials of Rablg Required per Total Body Weight of Client

Total Weight		# of Vials	Total Weight		# of Vials
$\leq 33 \text{ lbs}$	$\leq 15 \text{ Kg}$	1	$>165 - 198 \text{ lbs}$	$>75 - 90 \text{ Kg}$	6
$>33 - 66 \text{ lbs}$	$>15 - 30 \text{ Kg}$	2	$>198 - 231 \text{ lbs}$	$>90 - 105 \text{ Kg}$	7
$>66 - 99 \text{ lbs}$	$>30 - 45 \text{ Kg}$	3	$>231 - 264 \text{ lbs}$	$>105 - 120 \text{ Kg}$	8
$>99 - 132 \text{ lbs}$	$>45 - 60 \text{ Kg}$	4	$>264 - 297 \text{ lbs}$	$>120 - 135 \text{ Kg}$	9
$>132 - 165 \text{ lbs}$	$>60 - 75 \text{ Kg}$	5	$>297 - 330 \text{ lbs}$	$>135 - 150 \text{ Kg}$	10

Note that the amount of Rablg administered may include administration of only a portion of one of the vials ordered. For example, a patient that weighs 52.5 kg and requires 1050 IU of Rablg should only have 3.5 vials administered, rather than 4 full vials, with the remainder of the Rablg in the 4th vial being discarded.